

On the Upper Bounds of MDS Codes

Jiansheng Yang*, Yunying Zhang

Department of Mathematics, Shanghai University
Shanghai 200444, China

Abstract

Let $M_q(k)$ be the maximum length of MDS codes with parameters q, k . In this paper, the properties of $M_q(k)$ are studied, and some new upper bounds of $M_q(k)$ are obtained. Especially we obtain that $M_q(q-1) \leq q+2$ ($q \equiv 4 \pmod{6}$), $M_q(q-2) \leq q+1$ ($q \equiv 4 \pmod{6}$), $M_q(k) \leq q+k-3$ ($q=36(5s+1)$, $s \in N$ and $k=6, 7$).

Keywords: MDS codes; Hamming distance; codes equivalence; weight distribution

1 Introduction

Let C be an (n, q^k, d) code, if $d = n - k + 1$, then C is called a maximum distance separable (MDS) code. MDS codes are at the heart of combinatorics and finite geometries. In their book [9] MacWilliams and Sloane describe MDS codes as one of the most fascinating chapters in all of coding theory. These codes can be linear or non-linear. Very little is known about non-linear $(n, q^k, n - k + 1)$ MDS codes. R.H.Bruck, H.J.Ryser, R.Silverman and A.A.Bruen had proved some results on MDS codes [4,5,10] early. Recently, T.L.Alderson studies MDS codes extension. And he has obtained some important results[1,2,3].

In this paper, we assume that $A = \{0, 1, 2, \dots, q-1\}$ is a additive group(not necessary cyclic group). Denote the maximum number n of an $(n, q^k, n - k + 1)$ MDS code over A by $M_q(k)$. If A is a field, then denote the maximum

*Supported by Shanghai Leading Academic Discipline Project Project NumberS30104.
Email: yjsyjs@staff.shu.edu.cn

number n of a linear $(n, q^k, n - k + 1)$ MDS code over A by $m_q(k)$. The Main Conjecture of $m_q(k)$ is the following.

$$m_q(k) = \begin{cases} q + 2 & \text{for } k = 3 \text{ and } k = q - 1 \text{ both } q \text{ even,} \\ q + 1 & \text{in all other cases.} \end{cases}$$

The Main Conjecture has been proved in some cases, [7] gave us a good summarize. For $M_q(k)$, it is well known that $M_q(k) \leq q + k - 1$. In [5], A.A.Bruen and R.Silverman proved that:

Theorem 1.1 [5] (1) If C is a $(q + k - 1; k)$ -MDS code with $k \geq 3$ and $q > 2$, then 4 divides q .

(2) If C is a $(q + k - 1; k)$ -MDS code with $k > 3$ and $q > 2$, then 36 divides q .

In [2], T.L.Alderson proved that:

Theorem 1.2 [2] (1) If 36 does not divide q and $k \geq 4$, then a q -ary (n, k) -MDS code satisfies $n \leq q + k - 3$.

(2) If $q > 2$ and $q \equiv 2 \pmod{4}$ then no q -ary $(q + 1; 3)$ -MDS codes exist.

In [11], Wang proved that

Theorem 1.3 [11] $M_q(q - 1) \leq q + 1$ for q is odd.

In this paper, we use the generalized weight enumerator(see below) and combinatorial methods to study $M_q(k)$. Some new upper bounds of $M_q(k)$ are obtained. Especially we obtain that $M_q(q - 1) \leq q + 2$ ($q \equiv 4 \pmod{6}$), $M_q(q - 2) \leq q + 1$ ($q \equiv 4 \pmod{6}$), $M_q(k) \leq q + k - 3$ ($q = 36(5s + 1)$, $s \in \mathbb{N}$ and $k = 6, 7$).

For the convenient, we introduce some notations and results as following.

Let C and D be two codes of length n over A . If there exist n permutations $\pi_1, \dots, \pi_n$ of the q elements and a permutation σ of the n coordinate positions such that $(u_1, \dots, u_n) \in C$ iff $(\pi_1(u_{\sigma(1)}), \dots, \pi_n(u_{\sigma(n)})) \in D$, then we call C is equivalent to D . If C is equivalent to D , then C and D have the same Hamming distance. It is clear, if $c_0 \in A^n$, $D = c_0 + C = \{c_0 + \alpha \mid \alpha \in C\}$ is equivalent to C . Then we may always assume that code C contains the zero element $\mathbf{0} = (0, 0, \dots, 0, 0)$.

The generalized weight enumerator which is introduced by M.El-Khamy and R.J.McEliece[6] is called the partition weight enumerator(PWE). Suppose the coordinate set $N = \{1, 2, \dots, n\}$ is partitioned into p disjoint subsets $N_1, \dots, N_p$, with $|N_i| = n_i$, for $i = 1, \dots, p$. Denoting this partition by $\mathcal{T}$, the $\mathcal{T}$ -weight profile of an $v \in A^n$ is defined as $\mathcal{W}_{\mathcal{T}}(v) = (\omega_1, \dots, \omega_p)$, where ω_i is the Hamming weight of v restricted to N_i . Given a code C of

length n , and an $(n_1, \dots, n_p)$ partition $\mathcal{T}$ of the n coordinates of C , the $\mathcal{T}$ -weight enumerator of C is defined as following.

$$A^{\mathcal{T}}(\omega_1, \dots, \omega_p) = |\{c \in C : \mathcal{W}_{\mathcal{T}}(c) = (\omega_1, \dots, \omega_p)\}|.$$

Theorem 1.4 [6] For an (n, q^k, d) MDS code C which contains the zero element, the p -partition weight enumerator is given by

$$A^{\mathcal{T}}(\omega_1, \dots, \omega_p) = E(\omega) \frac{\binom{n_1}{\omega_1} \binom{n_2}{\omega_2} \cdots \binom{n_p}{\omega_p}}{\binom{n}{\omega}}$$

where $\omega = \sum_{i=1}^n \omega_i$, $E(\omega) = |\{c \in C : \mathcal{W}(c) = \omega\}|$ and $\mathcal{W}(c)$ is the Hamming weight of C .

Remark: In the proof, [6] assume that A is a field, however, the proof is true for any A (It only need the MDS codes have the zero element). Therefore, the formula holds for non-linear MDS codes which contain the zero element.

For an (n, q^k, d) MDS code over A , the weight distribution is known as

$$E(\omega) = (q-1) \binom{n}{\omega} \sum_{j=0}^{\omega-d} (-1)^j \binom{\omega-1}{j} q^{\omega-d-j}$$

where $\omega \geq d$ [9], and we can know that the formula holds for MDS codes (which contain the zero element) not only for linear MDS codes [8].

For any $\alpha = (a_1, \dots, a_n) \in A^n$, define the support of α by $Supp \alpha = \{i | a_i \neq 0, 1 \leq i \leq n\}$, and $\overline{Supp \alpha} = \{j | a_j = 0, 1 \leq j \leq n\}$.

2 New Upper Bounds for MDS Codes

Theorem 2.1 If $q \equiv 4 \pmod{6}$, then $M_q(q-1) \leq q+2$.

Proof: Suppose C is an $(q+3, q^{q-1}, 5)$ (q is even) MDS code which contains the zero element. The partition $\mathcal{T}$ is given as following.

$$\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2, \quad \mathcal{T}_1 = \{1, 2, 3\}, \quad \mathcal{T}_2 = \{4, 5, \dots, q+3\}.$$

$$n_1 = |\mathcal{T}_1| = 3, \quad n_2 = |\mathcal{T}_2| = q, \quad \omega_1 = 2, \quad \omega_2 = 3.$$

From Theorem 1.4, we have

$$A^{\mathcal{T}}(2, 3) = E(5) \frac{\binom{3}{2} \binom{q}{3}}{\binom{q+3}{5}} = \frac{3q(q-1)^2(q-2)}{6}.$$

where $E(5) = (q-1)\binom{q+3}{5}$. For (x, y) , there are altogether $(q-1)^2$ pairs (x, y) with $x, y \in S$ where $S = \{1, 2, \dots, q-1\}$. Thus there exists $(a, b) \in S$ such that

$$|C_{a,b,0}| \geq \frac{3q(q-1)^2(q-2)}{6 \times 3(q-1)^2} = \frac{q(q-2)}{6}.$$

where

$$C_{a,b,0} = \{(a_1, a_2, a_3, \dots, a_{q+3}) \in C | a_1 = a, a_2 = b, a_3 = 0, a_k \in A, k = 4, 5, \dots, q+3\}.$$

Since C is an MDS code, w.l.g. we may assume $a = 1, b = 1$. Then

$$C_{1,1,0} = \{(1, 1, 0, a_4, \dots, a_{q+3}) \in C | a_k \in A, k = 4, 5, \dots, q+3\}.$$

Let

$$C_i = C_{1,1,0,i} = \{(1, 1, 0, a_4, \dots, a_{q+3}) \in C | a_i \neq 0\} \ (i \in \{4, 5, \dots, q+3\}).$$

We will prove that $|C_{1,1,0}| = \frac{q(q-2)}{6}$ when q is even. If this is not true, we have $|C_{1,1,0}| > \frac{q(q-2)}{6}$, then there exists i , w.l.g. assume $i=4$, such that

$$|C_4| \geq \frac{|C_{1,1,0}| \binom{3}{1}}{\binom{q}{1}} > \frac{3q(q-2)}{6q} = \frac{q-2}{2}.$$

Assume $\alpha = (1, 1, 0, a_4, \dots, a_{q+3})$, $\beta = (1, 1, 0, b_4, \dots, b_{q+3}) \in C_4$, we have $a_4 \neq 0$, $b_4 \neq 0$. If $i \in \text{Supp}\alpha \cap \text{Supp}\beta$ ($5 \leq i \leq q+3$), then we have $d(\alpha, \beta) \leq 4$, a contradiction. Thus we have $\text{Supp}\alpha \cap \text{Supp}\beta = \{1, 2, 4\}$. Let $\alpha_1, \alpha_2, \dots, \alpha_t \in C_4$. Since $\text{Supp}\alpha_i \cap \text{Supp}\alpha_j = \{1, 2, 4\}$ for all $i \neq j$ and $\omega(\alpha) = 5$, we have $\cup_{i=1}^t |\text{Supp}\alpha_i| = 2t + 3$. This implies $2t + 3 \leq q + 2$, i.e. $t \leq \frac{q-1}{2}$. Since q is even, we have $t \leq \frac{q-2}{2}$.

Hence $|C_4| \leq \frac{q-2}{2}$ (q is even), a contradiction.

By this, we have

$$|C_{1,1,0}| = \frac{q(q-2)}{6} \ (q \text{ is even}).$$

Thus $\frac{q(q-2)}{6}$ must be an integer, however, if $q \equiv 4 \pmod{6}$, $\frac{q(q-2)}{6}$ is not an integer. Therefore, if $q \equiv 4 \pmod{6}$, then $M_q(q-1) \leq q+2$. $\square$

Theorem 2.2 If q is even and $(l+2)!$ does not divide $(q+l-1) \cdots (q+1)q(q-2)$ where $l \geq 1$, then $M_q(q-2) \leq q+l$.

Proof: Suppose C is an $(q + l + 1, q^{q-2}, l + 4)$ (q is even) MDS code which contains the zero element. The partition $\mathcal{T}$ is given as following.

$$\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2, \quad \mathcal{T}_1 = \{1, 2\}, \quad \mathcal{T}_2 = \{3, 4, \dots, q + l + 1\}.$$

$$n_1 = |\mathcal{T}_1| = 2, \quad n_2 = |\mathcal{T}_2| = q + l - 1, \quad \omega_1 = 2, \quad \omega_2 = l + 2.$$

From Theorem 1.4, we have

$$A^{\mathcal{T}}(2, l + 2) = E(l + 4) \frac{\binom{2}{2} \binom{q+l-1}{l+2}}{\binom{q+l+1}{l+4}} = \binom{q+l-1}{l+2} (q - 1).$$

where $E(l + 4) = (q - 1) \binom{q+l+1}{l+4}$. For (x, y) , there are altogether $(q - 1)^2$ pairs (x, y) with $x, y \in S$ where $S = \{1, 2, \dots, q - 1\}$. Thus there exists $(a, b) \in S$ such that

$$|C_{a,b}| \geq \frac{\binom{q+l-1}{l+2} (q - 1)}{(q - 1)^2} = \frac{\binom{q+l-1}{l+2}}{q - 1}.$$

where $C_{a,b} = \{(a_1, a_2, a_3, \dots, a_{q+l+1}) | a_1 = a, a_2 = b, a_k \in A, k = 3, 4, \dots, q + l + 1\}$.

Since C is an MDS code, we may assume $a = 1, b = 1$. Then

$$C_{1,1} = \{(1, 1, a_3, \dots, a_{q+l+1}) | a_k \in A, k = 3, 4, \dots, q + l + 1\}.$$

Let $B_{\underbrace{i, j, \dots, k}_l} = C_{1,1, \underbrace{i, j, \dots, k}_l} = \{(1, 1, a_3, \dots, a_{q+l+1}) | a_k \in A, k = 3, 4, \dots, q + l + 1 \text{ and } a_i, a_j, \dots, a_k \neq 0\}$, where $i, j, \dots, k$ are the l distinct numbers of $\{3, 4, \dots, q + l + 1\}$.

We claim that $|C_{1,1}| = \frac{\binom{q+l-1}{l+2}}{q-1}$ when q is even. If this is not true, since $|C_{1,1}| \geq \frac{\binom{q+l-1}{l+2}}{q-1}$, we have $|C_{1,1}| > \frac{\binom{q+l-1}{l+2}}{q-1}$ and there exist $i, j, \dots, k$, w.l.g. assume the l numbers are $3, 4, \dots, l + 2$, such that

$$|B_{3,4,\dots,l+2}| \geq \frac{|C_{1,1}| \binom{l+2}{l}}{\binom{q+l-1}{l}} > \frac{q-2}{2}.$$

Assume $\alpha = (1, 1, a_3, \dots, a_{q+l+1})$, $\beta = (1, 1, b_3, \dots, b_{q+l+1}) \in B_{3,4,\dots,l+2}$, we have $a_r \neq 0$, $b_s \neq 0$, $3 \leq r, s \leq l + 2$. If $i \in \text{Supp}\alpha \cap \text{Supp}\beta$ ($l + 3 \leq i \leq q + l + 1$), since $|\text{Supp}\alpha| = |\text{Supp}\beta| = l + 4$, then we have $d(\alpha, \beta) \leq l + 3$, a contradiction. Thus we have $\text{Supp}\alpha \cap \text{Supp}\beta = \{1, 2, \dots, l + 2\}$. Let $\alpha_1, \alpha_2, \dots, \alpha_t \in B_{3,4,\dots,l+2}$. Since $\text{Supp}\alpha_i \cap \text{Supp}\alpha_j = \{1, 2, \dots, l + 2\}$ for all

$i \neq j$ and $\omega(\alpha) = l + 4$, we have $\cup_{i=1}^t |Supp \alpha_i| = 2t + l + 2$. This implies $2t + l + 2 \leq q + l + 1$, i.e. $t \leq \frac{q-1}{2}$. Since q is even, we have $t \leq \frac{q-2}{2}$.

Hence $|B_{3,4,\dots,l+2}| \leq \frac{q-2}{2}$ (q is even), a contradiction.

By this, we have

$$|C_{1,1}| = \frac{\binom{q+l-1}{l+2}}{q-1} = \frac{(q+l-1) \cdots (q+1)q(q-2)}{(l+2)!} \quad (q \text{ is even}).$$

Thus $\frac{(q+l-1) \cdots (q+1)q(q-2)}{(l+2)!}$ must be an integer. Therefore, if q is even and $(l+2)!$ does not divide $(q+l-1) \cdots (q+1)q(q-2)$ where $l \geq 1$, then $M_q(q-2) \leq q+l$. $\square$

By calculating, we can get the following.

Corollary 2.2.1 $M_q(q-2) \leq q+1$ ($q \equiv 4 \pmod{6}$).

Corollary 2.2.2 $M_q(q-2) \leq q+3$ ($q \equiv 6 \text{ or } 26 \pmod{30}$).

Corollary 2.2.3 $M_q(q-2) \leq q+5$ ($q \equiv 8 \text{ or } 36 \pmod{42}$).

Theorem 2.3 If q is even and $(k-1)!$ does not divide $(q+k-4) \cdots (q+1)q(q-2)$ where $k \geq 4$, then $M_q(k) \leq q+k-3$.

Proof: Suppose C is an $(q+k-2, q^k, q-1)$ (q is even) MDS code which contains the zero element. The partition $\mathcal{T}$ is given as following.

$$\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2, \quad \mathcal{T}_1 = \{1, 2\}, \quad \mathcal{T}_2 = \{3, 4, \dots, q+k-2\}.$$

$$n_1 = |\mathcal{T}_1| = 2, \quad n_2 = |\mathcal{T}_2| = q+k-4, \quad \omega_1 = 2, \quad \omega_2 = q-3.$$

From Theorem 1.4, we have

$$A^{\mathcal{T}}(2, q-3) = E(q-1) \frac{\binom{2}{2} \binom{q+k-4}{q-3}}{\binom{q+k-2}{q-1}} = \binom{q+k-4}{q-3} (q-1).$$

where $E(q-1) = (q-1) \binom{q+k-2}{q-1}$. For (x, y) , there are altogether $(q-1)^2$ pairs (x, y) with $x, y \in S$ where $S = \{1, 2, \dots, q-1\}$. Thus there exists $(a, b) \in S$ such that

$$|C_{a,b}| \geq \frac{\binom{q+k-4}{q-3} (q-1)}{(q-1)^2} = \frac{\binom{q+k-4}{q-3}}{q-1}. \quad (1)$$

where $C_{a,b} = \{(a_1, a_2, a_3, \dots, a_{q+k-2}) | a_1 = a, a_2 = b, a_m \in A, m = 3, 4, \dots, q+k-2\}$.

Since C is an MDS code, we may assume $a = 1, b = 1$. Then

$$C_{1,1} = \{(1, 1, a_3, \dots, a_{q+k-2}) | a_m \in A, m = 3, 4, \dots, q+k-2\}.$$

Let $B_{\underbrace{i, j, \dots, r}_{k-3}} = C_{1,1, \underbrace{i, j, \dots, r}_{k-3}} = \{(1, 1, a_3, \dots, a_{q+k-2}) | a_m \in A, m = 3, 4, \dots, q+k-2 \text{ and } a_i = a_j = \dots = a_r = 0\}$, where $i, j, \dots, r$ are the $k-3$ distinct numbers of $\{3, 4, \dots, q+k-2\}$.

We will prove that $|C_{1,1}| = \frac{\binom{q+k-4}{q-3}}{q-1}$ when q is even. If this is not true, we have $|C_{1,1}| > \frac{\binom{q+k-4}{q-3}}{q-1}$ and there exist $i, j, \dots, r$, w.l.g. assume the $k-3$ numbers are $3, 4, \dots, k-1$, such that

$$|B_{3,4,\dots,k-1}| \geq \frac{|C_{1,1}| \binom{k-1}{k-3}}{\binom{q+k-4}{k-3}} > \frac{q-2}{2}.$$

Assume $\alpha = (1, 1, a_3, \dots, a_{q+k-2})$, $\beta = (1, 1, b_3, \dots, b_{q+k-2}) \in B_{3,4,\dots,k-1}$, we have $a_r = 0$, $b_s = 0, 3 \leq r, s \leq k-1$. If $i \in \overline{Supp\alpha} \cap \overline{Supp\beta}$ ($k \leq i \leq q+k-2$), then we have $d(\alpha, \beta) \leq q-2$, a contradiction. Thus we have $\overline{Supp\alpha} \cap \overline{Supp\beta} = \{3, 4, \dots, k-1\}$. Let $\alpha_1, \alpha_2, \dots, \alpha_t \in B_{3,4,\dots,k-1}$. Since $\overline{Supp\alpha_i} \cap \overline{Supp\alpha_j} = \{3, 4, \dots, k-1\}$ for all $i \neq j$ and $\omega(\alpha) = q-1$, we have $\cup_{i=1}^t |\overline{Supp\alpha_i}| = 2t + k - 3$. This implies $2t + k - 3 \leq q + k - 4$, i.e. $t \leq \frac{q-1}{2}$. Since q is even, we have $t \leq \frac{q-2}{2}$.

Hence $|B_{3,4,\dots,k-1}| \leq \frac{q-2}{2}$ (q is even), a contradiction.

By this, we have

$$|C_{1,1}| = \frac{\binom{q+k-4}{q-3}}{q-1} = \frac{(q+k-4) \cdots (q+1)q(q-2)}{(k-1)!} \quad (q \text{ is even})$$

Thus $\frac{(q+k-4) \cdots (q+1)q(q-2)}{(k-1)!}$ must be an integer. Therefore, if q is even and $(k-1)!$ does not divide $(q+k-4) \cdots (q+1)q(q-2)$ where $k \geq 4$, then $M_q(k) \leq q+k-3$. $\square$

By the Theorem 2.3, we have the following.

Corollary 2.3.1 $M_q(k) \leq q+k-3$ ($q = 36(5s+1)$, $s \in N$ and $k = 6, 7$).

3 Conclusion

In this paper, we use the generalized weight enumerator and combinatorial methods to study $M_q(k)$ which denote the maximum number n of an

$(n, q^k, n - k + 1)$ MDS code. Compared to Theorem 1.1, Theorem 1.2 and Theorem 1.3, we obtain some new upper bounds of $M_q(k)$. Especially we obtain that $M_q(q - 1) \leq q + 2(q \equiv 4(\text{mod } 6))$, $M_q(q - 2) \leq q + 1(q \equiv 4(\text{mod } 6))$, $M_q(k) \leq q + k - 3$ ($q = 36(5s + 1)$, $s \in \mathbb{N}$ and $k = 6, 7$).

References

- [1] T.L. Alderson, "On MDS codes and Bruen-Silverman codes," Ph.D. Thesis, *University of Western Ontario*, 2002.
- [2] T.L.Alderson, "Extending MDS codes," *Annals of Combinatorics*, vol. 9, pp. 125-135, 2005.
- [3] T. L. Alderson and A. A. Bruen, "Codes from cubic curves and their extensions," *the electronic journal of combinatorics*, vol. 15, 2008.
- [4] R.H.Bruck and H.J.Ryser, "The nonexistence of certain finite projective planes," *Canad J.Math*, vol. 1, pp. 88-93, 1949.
- [5] A.A.Bruen and R.Silverman, "On the nonexistence of certain MDS codes and projective planes," *Math.Z*, vol. 183, pp. 171-175, 1983.
- [6] M.El-Khamy and R.J.McEliece, "The Partition Weight Enumerator of MDS Codes and its Applications," *Information Theory*, vol. 9, pp. 926-930, 2005.
- [7] J. W. P. Hirschfeld, "The Main Conjecture for MDS Codes," Lecture Notes In Computer Science; Proceedings of the 5th IMA Conference on Cryptography and Coding, *Springer-Verlag London, UK*, vol. 1025, pp. 44-52, 1995.
- [8] Ludo M.G.M. and Tolhuizen, "On Maximum Distance Separable codes over alphabets of arbitrary size," *Information Theory*, vol. 7, pp. 431, 1994.
- [9] F.J.MacWilliams and N.J.A.Slane, "Theory of Error-Correcting Codes," *North-Holland, Amsterdam*, pp. 317-329, 1977.
- [10] R.Silverman, "A metrization for power-sets with applications to combinatorial analysis," *Canad.J.Math*, vol. 12, pp. 158-176, 1960.
- [11] D.X.Wang, "The MDS codes of small dimension and small weight," Ms.Thesis, *University of Shanghai*, 2008.